

Example 1

High scab risk / low liver fluke risk - scan ELISA negative.



www.scops.org.uk




MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY
1	2	3	4 100 Ewe lambs bought at market. Decide to test for sheep scab rather than treat immediately.	5 Treated for Roundworms Group 4- AD + mox used (silver option)	6	7 Turn out to Church Field where sheep have grazed until recently but now isolated from the rest of the flock.
<div style="background-color: #e67e22; color: white; padding: 5px; display: inline-block; border-radius: 10px;"> ➔ Housed on arrival for 48 hours </div>						
8	9	10	11	12	13	14
15	16	17	18	19 Take mob FEC sample to test the wormers have worked. Take blood samples for scab ELISA test	20	21
22	23	24 Results of scab test are negative.	25	26	27	28 Ewe lambs have completed 3 weeks in isolation and can now move to join the rest of the flock if required.
29	30	31				

Example 1

High scab risk / low liver fluke risk - scan ELISA negative.



www.scops.org.uk



MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY
1	2	3	4 100 Ewe lambs bought at market. Decide to test for sheep scab rather than treat immediately.	5 Treated for Roundworms Group 4- AD + mox used (silver option)	6	7 Turn out to Church Field where sheep have grazed until recently but now isolated from the rest of the flock.
Housed on arrival for 48 hours						
8	9	10	11	12	13	14
15	16	17	18	19 Take mob FEC sample to test the wormers have worked. Take blood samples for scab ELISA test	20	21
22	23	24 Results of scab test are negative.	25	26	27	28 Ewe lambs have completed 3 weeks in isolation and can now move to join the rest of the flock if required.
29	30	31				