

Example 2

Low scab risk / low liver fluke risk - scan ELISA negative.



www.scops.org.uk




MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY
1	2	3	4 5 shearling rams bought direct from a farm. Scab risk considered low	5 Treated for Roundworms (Group 4-AD + 5 SI used (gold))	6	7 Turn out to small paddock next to sheds where sheep have grazed until recently but now isolated from the rest of the flock.
<div style="background-color: #e67e22; color: white; padding: 5px; display: inline-block; border-radius: 10px;"> ➔ </div> Housed on arrival for 48 hours						
8	9	10	11	12	13	14
15	16	17	18	19 Take mob FEC sample to test the wormers have worked effectively and blood for the ELISA test for sheep scab.	20 FEC test shows no worm eggs so treatment was effective.	21
22	23	24 Results of scab test are negative.	25	26	27	28 Rams are now clear and can join others and/or go out to work with the ewes.
29	30	31				

Example 2

Low scab risk / low liver fluke risk - scan ELISA negative.



www.scops.org.uk



MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY
1	2	3	4 5 shearling rams bought direct from a farm. Scab risk considered low	5 Treated for Roundworms (Group 4-AD + 5 SI used (gold)	6	7 Turn out to small paddock next to sheds where sheep have grazed until recently but now isolated from the rest of the flock.
1 → 2 → 3						
8	9	10	11	12	13	14
3						
15	16	17	18	19 Take mob FEC sample to test the wormers have worked effectively and blood for the ELISA test for sheep scab.	20 FEC test shows no worm eggs so treatment was effective.	21
4						
22	23	24 Results of scab test are negative.	25	26	27	28 Rams are now clear and can join others and/or go out to work with the ewes.
5						
29	30	31				